


Introductory Chemistry
Chem 103

Chapter 9 – Covalent Bonding and Molecules

Lecture Slides



Covalent Molecules


Covalent bonds:

Two atoms share electrons
Occur between nonmetal atoms



Octet rule:


Atoms are stabilized by having 8 electrons in the valence shell



Lewis structures:


Show the arrangement of covalently bonded atoms

Use a dash to represent two shared electrons




Covalent Double and Triple Bonds


Covalent double bonds:




Covalent triple bonds:




Pairs of Electrons in Compounds



In Most Molecules, Atoms Follow the Octet Rule




**In Most Molecules, Atoms Follow the Octet Rule,
Continued**




Exceptions to the Octet Rule


incomplete octet



expanded octet



Complete octet



Expanded octet

	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95
		34 Se 78.97	35 Br 79.90	36 Kr 83.80
			53 I 126.90	54 Xe 131.29
				86 Rn (222)

Drawing Lewis Structures



1. Add up all the valence electrons.
2. Frame the structure.
3. Fill octets on outer atoms first.
4. Fill the octet on the central atom.
 - any remaining electrons on central atom
 - use double/triple bonds if needed

1A	2A	Main group number						3A	4A	5A	6A	7A	8A
1	2	Valence electrons						3	4	5	6	7	8

Revell, Introductory Chemistry, 2e, © 2021
W. H. Freeman and Company

Drawing Lewis Structures Practice

Draw a Lewis structure for nitrogen trichloride, NCl_3 .





1. Sum electrons.
2. Draw framework.
3. Fill octets on outer atoms.
4. Fill octet on central atom.

5A	6A	7A	2
15	16	17	He
7	8	9	10
N	O	F	Ne
14.01	16.00	19.00	20.18
15	16	17	18
P	S	Cl	Ar
30.97	32.06	35.45	39.95

Drawing Lewis Structures, More Practice

Formaldehyde, CH_2O , is commonly used to manufacture plastics.
Draw the Lewis structure for a formaldehyde molecule.




1. Sum electrons.
2. Draw framework.
3. Fill octets on outer atoms.
4. Fill octet on central atom.

4A	5A	6A	7A	2
14	15	16	17	He
6	7	8	9	10
C	N	O	F	Ne
12.01	14.01	16.00	19.00	20.18
14	15	16	17	18

Molecules and Charge


polyatomic ions groups of atoms with an overall charge

formal charges a method of identifying charged sites





Revel, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

One Electron From Each Bond is Assigned to an Atom









Calculating Formal Charge Practice

$$\text{Formal charge} = \text{Valence electrons in the neutral atom} - \text{Number of covalent bonds} - \text{Number of unshared electrons}$$




Oxygen and Nitrogen Atoms Often Have Formal Charges



Calculating Formal Charge, More Practice

Automotive air bags contain sodium azide, NaN_3 . The Lewis structure for the azide ion (without charges) is shown. Calculate the formal charge on each atom in this structure. What is the overall charge of the azide ion?



Drawing Lewis Structures for Polyatomic Ions


- Similar to neutral molecules
- Consider charge when finding the number of valence electrons

How many valence electrons are in a hydroxide ion?




Lewis Structures for Polyatomic Ions Practice

Draw a Lewis structure for the nitrite ion, NO_2^- . Show all nonzero formal charges.



= 18 valence electrons



1. Sum electrons.
2. Draw framework.
3. Fill octets on outer atoms.
4. Fill octet on central atom.




	5A	6A	7A	2
	15	16	17	He
7	N	O	F	10
14.01	16.00	19.00	20.18	Ne
15	P	S	Cl	18
30.97	32.06	35.45	39.95	Ar

Identifying the Best Lewis Structure




What is the best structure for phosgene, COCl_2 ?



or





Resonance Structures



Revel, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Resonance Structures, Continued


- a set of structures that show how electrons are distributed.
- used when a single Lewis structure is insufficient.



Revel, Introductory C

Ions With Resonance Structures Spread Charges Over Multiple Atoms


NO_3^-	SO_4^{2-}
NO_2^-	CO_3^{2-}
PO_4^{3-}	$\text{C}_2\text{H}_3\text{O}_2^-$



Only 2nd bonds and lone pairs change in resonance structures.

Using Resonance Structures to Calculate Formal Charge


The nitrate ion (NO_3^-) has three major resonance structures. Draw each structure. Based on these structures, what is the average charge on each oxygen atom?



charge on each oxygen: $-\frac{2}{3}$

Shapes of Molecules

CO₂
gas



H₂O
liquid






Brilliant Eye/Shutterstock


Predicting Molecular Shapes

Valence Shell Electron Pair Repulsion


VSEPR



Predicting Molecular Shapes, Continued




Electronic geometry
Arrangement of electrons
around the central atom




Molecular geometry
Shape caused by the
arrangement of atoms

Revell, *Introductory Chemistry*, 2e, © 2021
W. H. Freeman and Company

Two Electron Sets: Linear



Electronic geometry
Linear




Molecular geometry
Linear

Revell, *Introductory Chemistry*, 2e,
© 2021 W. H. Freeman and Company


double and triple
bonds count as 1 “set”

Geometric Stability


Which geometry is most stable?



a.




b.




c.


2 sets of electrons: Linear



Three Electron Sets: Trigonal Planar




Electronic geometry
Trigonal planar




Molecular geometry
Trigonal planar

Revell, *Introductory Chemistry*, 2e, © 2021 W. H. Freeman and Company

Three Electron Sets: Trigonal Planar, Continued





Electronic geometry
Trigonal planar



Molecular geometry
Bent


Revell, *Introductory Chemistry*, 2e, © 2021
W. H. Freeman and Company

Four Electron Sets: Tetrahedral, Part 1



In an X-shape, the bonds are all 90° apart.


However, if we rotate two of the bonds, we create more space...



The tetrahedron is like two V-shapes in planes that are perpendicular to each other.


Revell, *Introductory Chemistry*, 2e, © 2021
W. H. Freeman and Company

Four Electron Sets: Tetrahedral, Part 2




Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Electronic geometry
Tetrahedral




Molecular geometry
Tetrahedral

Four Electron Sets: Tetrahedral, Part 3



Electronic geometry
Tetrahedral




Molecular geometry
Trigonal pyramidal

Revell, Introductory Chemistry, 2e, © 2021
W. H. Freeman and Company

Four Electron Sets: Tetrahedral, Part 4






Electronic geometry
Tetrahedral




Molecular geometry
Bent

Revell, *Introductory Chemistry*, 2e,
© 2021 W. H. Freeman and Company

Electronic and Molecular Geometry

Electron sets	Electronic geometry	Model	Bonding sets	Lone pairs	Molecular geometry	Examples
2	Linear		2	0	Linear	$\text{:}\ddot{\text{O}}=\text{C}=\ddot{\text{O}}\text{:}$
3	Trigonal Planar		3	0	Trigonal Planar	$\text{H}-\text{C}(\text{O})-\text{H}$
			2	1	Bent	$\text{:}\ddot{\text{O}}-\text{N}(\text{O})-\ddot{\text{O}}\text{:}^-$
4	Tetrahedral		4	0	Tetrahedral	$\text{H}-\text{C}(\text{H})_3$
			3	1	Trigonal pyramidal	$\text{H}-\text{N}(\text{H})_3$
			2	2	Bent	$\text{H}-\text{O}(\text{H})_2$

Electronic and Molecular Geometry Practice




Trigonal planar

Tetrahedral

Electronic and Molecular Geometry, More Practice

Lithium carbonate is a simple ionic compound that is widely used to treat bipolar disorder. What is the molecular geometry of the carbonate ion?




*Electronic geometry:
Trigonal planar*

*Molecular geometry:
Trigonal planar*

Polar Bonds and Molecules

Polar covalent bond atoms do not share the electrons evenly







Electronegativity

how strongly atoms pull bonded electrons

																		H 2.1																				
Li 1.0	Be 1.5											B 2.0	C 2.5	N 3.0	O 3.5	F 4.0																						
Na 0.9	Mg 1.2											Al 1.5	Si 1.8	P 2.1	S 2.5	Cl 3.0																						
K 0.8	Ca 1.0	Sc 1.3	Ti 1.5	V 1.6	Cr 1.6	Mn 1.5	Fe 1.8	Co 1.8	Ni 1.8	Cu 1.9	Zn 1.7	Ga 1.6	Ge 1.8	As 2.0	Se 2.4	Br 2.8																						
R 0.8	Sr 1.0	Y 1.2	Zr 1.4	Nb 1.6	Mo 1.8	Tc 1.9	Ru 2.2	Rh 2.2	Pd 2.2	Ag 1.9	Cd 1.7	In 1.7	Sn 1.8	Sb 1.9	Te 2.2	I 2.5																						
Cs 0.7	Ba 0.9	La 1.1	Hf 1.3	Ta 1.5	W 1.7	Re 1.9	Os 2.2	Ir 2.2	Pt 2.2	Au 2.4	Hg 1.9	Tl 1.8	Pb 1.8	Bi 1.9	Po 2.0	At 2.2																						
Fr 0.7	Ra 0.9																																					

Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Comparing Covalent, Polar Covalent and Ionic Bonds



Revell, Introductory
Chemistry, 2e, © 2021
W. H. Freeman and
Company

Covalent: < 0.5


Polar Covalent: $0.5 - 2.0$

Ionic: > 2.0

																H 2.1												
Li 1.0		Be 1.5																B 2.0		C 2.5		N 3.0		O 3.5		F 4.0		
Na 0.9		Mg 1.2																Al 1.5		Si 1.8		P 2.1		S 2.5		Cl 3.0		
K 0.8		Ca 1.0		Sc 1.3	Ti 1.5	V 1.6	Cr 1.6	Mn 1.5	Fe 1.8	Co 1.8	Ni 1.8	Cu 1.9	Zn 1.7	Ga 1.6	Ge 1.8	As 2.0	Se 2.4	Br 2.8										
R 0.8		Sr 1.0		Y 1.2	Zr 1.4	Nb 1.6	Mo 1.8	Tc 1.9	Ru 2.2	Rh 2.2	Pd 2.2	Ag 1.9	Cd 1.7	In 1.7	Sn 1.8	Sb 1.9	Te 2.2	I 2.5										
Cs 0.7		Ba 0.9		La 1.1	Hf 1.3	Ta 1.5	W 1.7	Re 1.9	Os 2.2	Ir 2.2	Pt 2.2	Au 2.4	Hg 1.9	Tl 1.8	Pb 1.8	Bi 1.9	Po 2.0	At 2.2										
Fr 0.9		Ra 0.9																										

Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

An Analogy for Polar Covalent Bonds



Kevin Revell

Classifying Bonds Practice

Which bond is more polar, a C-O bond or an F-S bond? Show the direction of polarity for both bonds.

C: 2.5

F: 4.0

O: 3.5

S: 2.5

Difference = 1.0





Difference = 1.5

Polar Covalent

Polar Covalent

B	C	N	O	F
2.0	2.5	3.0	3.5	4.0
Al	Si	P	S	Cl
1.5	1.8	2.1	2.5	3.0

F-S is more polar




Molecules with dipoles

Molecular dipole an overall polarity in a molecule

net dipole

dipole




polar covalent bonds

shape




Rich Carey/Shutterstock

Identifying Molecules with a Net Dipole




Net Dipole



No Net Dipole


Identifying Molecules with a Net Dipole, Continued



Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company


Identifying Molecules with a Net Dipole Practice

Which of these have a net dipole?





Non-polar bonds

No dipole



Polar bonds

No net dipole



Polar bonds

Net dipole


H
2.1

B	C	N	O	F
2.0	2.5	3.0	3.5	4.0
Al	Si	P	S	Cl
1.5	1.8	2.1	2.5	3.0


How Dipoles Affect Properties – A Preview



Linear



No net dipole



Bent




Net dipole




Brilliant Eye/Shutterstock

How Dipoles Affect Properties – A Preview, Continued

H_2O
Net dipole
Liquid at 25 °C



CO_2
No net dipole
Gas at 25 °C




Revell, *Introductory Chemistry*, 2e, © 2021
W. H. Freeman and Company

Introductory Chemistry
Chem 103

Chapter 10 – Solids, Liquids, Gases

Lecture Slides




Interactions between Particles

	Atomic/Molecular Arrangement	Macroscopic Properties
Solid	Particles are close together and held in a fixed place.	Definite shape and volume
Liquid	Particles are close together but move freely past each other.	Definite volume; Adopts the shape of the container.
Gas	Particles are far apart and have very little interaction.	Adopts shape and volume of container

Phase change

A transition from one state of matter to another.



Ondrej Prosky/Shutterstock
 Revell, *Introductory Chemistry*, 2e, © 2021 W. H. Freeman and Company

The Forces Between Particles Influence Physical Properties

Stronger forces between particles - higher melting and boiling points.




Solids and Liquids



Ionic Substances

Lattices: rigid frameworks of atoms, molecules or ions.




Revell, *Introductory Chemistry*, 2e, © 2021 W. H. Freeman and Company

Compound	Melting Point (°C)
NaCl	801
KCl	770
MgCl ₂	714
CaO	2,572
Al ₂ O ₃	2,072

Metallic Substances

- Form lattices of tightly packed atoms.
- Electrons move easily between atoms.
- Shapes of metals are easily altered.
 - Malleable
 - Ductile



Love Silhouette/Shutterstock
Revell, *Introductory Chemistry*, 2e, © 2021 W. H. Freeman and Company
Photo credit: Love Silhouette/Shutterstock

Metallic Substances, Continued




Element	Melting Point (°C)
Lead	327
Aluminum	660
Gold	1,064
Copper	1,085
Iron	1,538

Molecular Substances

Forces within molecules: covalent bonds


Forces between molecules: intermolecular forces



Revell, *Introductory Chemistry*,
2e, © 2021 W. H. Freeman and
Company

covalent networks: lattices of covalent bonds that form giant molecules

covalent networks: lattices of covalent bonds that form giant molecules




polymers: contain long chains of covalently-bonded atoms

polymers: contain long chains of covalently-bonded atoms




Caia Images/Superstock
Revell, *Introductory Chemistry*, 2e, © 2021 W. H. Freeman and Company

Solids and Liquids Summary




- Ionic Substances


- Metallic Substances



- Molecular Substances




- Covalent Networks and Polymers



Forces Within and Between Molecules

Forces within molecules: covalent bonds

Forces between molecules: intermolecular forces



Revel, Introductory Chemistry,
2e, © 2021 W. H. Freeman and
Company


Forces Between Molecules

intermolecular forces

1. Dipole-dipole Interactions
2. Hydrogen bonds
3. Dispersion forces

Dipole-Dipole Interactions, Part 1

Attractions between polar covalent molecules:



Revell, *Introductory Chemistry*, 2e, © 2021 W. H. Freeman and Company




NikolayN/Shutterstock

Photo credit: NikolayN/Shutterstock

Dipole-Dipole Interactions, Part 2

Dipole
Higher Melting Point
Higher Boiling Point





Revel, Introductory Chemistry, 2e, © 2011 W. H. Freeman and Company

No dipole

Dipole-Dipole Interactions, Part 3

	$\ddot{\text{O}}=\text{C}=\ddot{\text{O}}$ Carbon dioxide	$\ddot{\text{O}}=\ddot{\text{S}}=\ddot{\text{O}}$ Sulfur dioxide	$\text{H}-\text{C}(\text{H})_2-\text{C}\equiv\text{N}:$ Acetonitrile
Geometry	Linear	Bent	Linear
Dipole	Zero	Small	Large
Boiling Point			


Hydrogen Bonding, Part 1

		$\text{H}-\text{C}\equiv\text{N:}$	
	Methane	Hydrogen cyanide	Water
Formula mass	16.0 u	27.0 u	18.0 u
Dipole strength*	0	2.98	1.85
Boiling point			

*These numbers convey the relative size of each dipole.

Hydrogen Bonding, Part 2


A strong intermolecular force between molecules containing H-F, H-O, or H-N bonds.





Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Hydrogen Bonding, Part 3

A strong intermolecular force between molecules containing H-F, H-O, or H-N bonds.



Hydrogen Bonds Explain the Properties of Water



Revell, *Introductory Chemistry*, 2e, © 2021
W. H. Freeman and Company



a



b




c

a: Hermann Bierbaiss/Science Source; b: o. vancay/depositphotos.com; c: © 1991 Richard Megna/Fundamental Photographs, NYC


London Dispersion Forces, Part 1

Weak intermolecular forces that result from instantaneous dipoles



London Dispersion Forces, Part 2

Weak intermolecular forces that result from instantaneous dipoles



Summary of Intermolecular Forces



Left: chair01/
Shutterstock; center:
Denis Kapekhiu/
Shutterstock; right:
Manuel Ploetz/
Shutterstock


Type	Description	Strength
Hydrogen bonding	molecules with H-F, H-O, or H-N bonds	strongest
Dipole-dipole forces	molecules with net dipole	
London dispersion forces	all covalent molecules	weakest

Describing Gases

particles are spaced far apart
very little interaction between particles




Philip and Karen Smith/Getty Images



Reuell, Introductory Chemistry, 2e, © 2021
W. H. Freeman and Company

Ideal Gas

1. Volume of particles is much less than container.
2. Particles have no attraction for each other.




Revel, Introductory Chemistry, 2e,
© 2021 W. H. Freeman and Company

Temperature
Volume
Pressure

Pressure

The force that gases exert on their surroundings.

$$\text{Pressure} = \frac{\text{Force}}{\text{Area}}$$



Revel, Introductory Chemistry, 2e,
© 2021 W. H. Freeman and Company




Cathyrose Melloan/Alamy

Measuring Pressure



SimpleFoto/Deposit Photos



www.denisowald.de/Getty Images



Top: PhotoLibrary/Getty Images; Bottom Left: Carolyn Frankel/Getty Images; Bottom Right: Mark Hunt/Design Pics Media/Getty Images

Measuring Pressure – Barometers

Barometer: a device used to measure atmospheric pressure

Millimeters of mercury (mm Hg)


1 mm Hg = 1 torr

Average air pressure at sea level:

760 mm Hg


760 torr

Standard pressure



Revel, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Measuring Pressure – Barometers Continued



Revel, *Introductory Chemistry*, 2e, © 2021 W. H. Freeman and Company



© Kianthoon/depositphotos.com

Measuring Pressure – Gauge Pressure

Gauge Pressure: The difference between the compressed gas pressure and the atmospheric pressure.



a



b



c



d

(a) Fotostock/Shutterstock; (b) Sergey Zagorodny/Shutterstock; (c) Dmitry Kalinovsky/Shutterstock; (d) Science Source

Measuring Pressure – Conversion Factors

1 **atmosphere** (atm) = 760 mm Hg (torr)

1 atm = 14.70 pounds per square inch (psi)

1 atm = 101.3 kilopascals (kPa)


1 atm = 1.013 bar

The Gas Laws, Part 1

- Boyle's Law
- Charles's Law
- Combined Gas Law



Philip and Karen Smith/Getty Images



Revel, Introductory Chemistry, 2e, © 2021
W. H. Freeman and Company

Boyle's Law

The pressure and volume of a gas are inversely related.

$$P \uparrow \quad V \downarrow$$

$$PV = \text{constant}$$

$$P_1 V_1 = P_2 V_2$$



Boyle's Law Practice

A commercial compressor stores 2.8 liters of air at a pressure of 150 psi. If this air is allowed to expand until the pressure is equal to 15 psi (just over atmospheric pressure), what volume will the air occupy?

$$P_1 V_1 = P_2 V_2$$

$$P_1 = 150 \text{ psi}$$

$$V_1 = 2.8 \text{ L}$$

$$P_2 = 15 \text{ psi}$$

$$V_2 = ?$$

$$V_2 = \frac{P_1 V_1}{P_2}$$

$$= \frac{(150 \text{ psi})(2.8 \text{ L})}{(15 \text{ psi})} = 28 \text{ L}$$

Charles's Law


At constant pressure, the volume of a gas is directly proportional to its temperature.

$$T \uparrow \quad V \uparrow$$

$$V \propto T$$

$$\frac{V}{T} = \text{constant}$$

Using Charles's Law to Find Absolute Zero



The Kelvin Scale

Absolute zero

-273.15 °C

0 K

$$\text{kelvin} = ^\circ\text{C} + 273.15$$

Working to the nearest degree:

$$\text{kelvin} = ^\circ\text{C} + 273$$

Solving Problems with Charles's Law

$$\frac{V}{T} = \text{constant}$$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

Charles's Law Practice

A balloon has a volume of 3.2 liters at room temperature (25 °C). The gas inside the balloon is then heated to 100 °C. What is the new volume of the balloon?

$$V_1 = 3.2 \text{ L}$$

$$V_2 = ?$$

$$T_1 = 25^\circ\text{C} + 273 = 298 \text{ K}$$

$$T_2 = 100^\circ\text{C} + 273 = 373 \text{ K}$$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$V_2 = \frac{V_1 T_2}{T_1}$$

$$= \frac{(3.2 \text{ L})(373 \text{ K})}{(298 \text{ K})} = 4.0 \text{ L}$$

The Combined Gas Law



$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

Photo credit: Carolyn Franks/Alamy

The Combined Gas Law Practice

A gas with a temperature of 280 K, a pressure of 200 kPa, and a volume of 25.8 L is compressed to 15.8 L, causing the pressure to increase to 350 kPa. What is the temperature of the gas under the new conditions?

$$P_1 = 200 \text{ kPa}$$

$$V_1 = 25.8 \text{ L}$$

$$T_1 = 280 \text{ K}$$

$$P_2 = 350 \text{ kPa}$$

$$V_2 = 15.8 \text{ L}$$

$$T_2 = ?$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$T_2 = \frac{P_2 V_2 T_1}{P_1 V_1}$$


$$= \frac{(350 \text{ kPa})(15.8 \text{ L})(280 \text{ K})}{(200 \text{ kPa})(25.8 \text{ L})}$$

$$= 300 \text{ K}$$

The Gas Laws, Part 2

- Boyle's Law
- Charles's Law
- Combined Gas Law

- Avogadro's Law
- Ideal Gas Law



Philip and Karen Smith/Getty Images

Avogadro's Law

If temperature and pressure are constant, the volume of a gas is proportional to the number of moles of gas present.

$$V \propto n$$

at Standard Temperature and Pressure (STP)...

$$T = 0\text{ }^{\circ}\text{C} \text{ (273 K)}$$

$$P = 1.0\text{ atm}$$

...1 mole of gas occupies 22.4 Liters



Shutterstock Images

The Ideal Gas Law

$$PV = nRT$$

- $R = 0.0821\text{ L}\cdot\text{atm}/\text{mol}\cdot\text{K}$
- T must be in kelvins
- P , V units must match gas constant

The Ideal Gas Law Practice

What volume does 1.00 mole of gas occupy at a temperature of 0.00 °C and a pressure of 1.00 atmospheres?

$$PV = nRT$$

$$T = 0.00\text{ }^{\circ}\text{C}$$

$$= 273.15\text{ K}$$

$$V = \frac{nRT}{P}$$

$$= \frac{(1.00\text{ mol})(0.0821\text{ L}\cdot\text{atm/mol}\cdot\text{K})(273.15)}{1.00\text{ atm}}$$

$$= 22.4\text{ L}$$



Shutterstock/Getty Images


Mixtures of Gases: Partial Pressure

partial pressure: The pressure caused by one gas in a mixture.

Adding up all partial pressures gives the total pressure.



Philip and Karen Smith/Getty Images



Revel, Investigating Chemistry, 2e, © 2017
W. H. Freeman and Company

Air:
78% nitrogen
21% oxygen

Partial Pressure Practice

If a 40.0-L cylinder is filled with 5.00 moles of nitrogen, 2.00 moles of oxygen, and 3.00 moles of carbon dioxide at a temperature of 400 K, what is the pressure inside the cylinder?

$$P_{N_2} = \frac{nRT}{V} = \frac{(5.00 \text{ mol})(0.0821 \text{ L}\cdot\text{atm/mol}\cdot\text{K})(400 \text{ K})}{40.0 \text{ L}} = 4.11 \text{ atm}$$

$$P_{O_2} = \frac{nRT}{V} = \frac{(2.00 \text{ mol})(0.0821 \text{ L}\cdot\text{atm/mol}\cdot\text{K})(400 \text{ K})}{40.0 \text{ L}} = 1.64 \text{ atm}$$

$$P_{CO_2} = \frac{nRT}{V} = \frac{(3.00 \text{ mol})(0.0821 \text{ L}\cdot\text{atm/mol}\cdot\text{K})(400 \text{ K})}{40.0 \text{ L}} = 2.46 \text{ atm}$$

$$\begin{aligned} P_{Total} &= P_{N_2} + P_{O_2} + P_{CO_2} \\ &= 4.11 \text{ atm} + 1.64 \text{ atm} + 2.46 \text{ atm} = 8.21 \text{ atm} \end{aligned}$$


Partial Pressure, More Practice

If a 40.0-L cylinder is filled with 5.00 moles of nitrogen, 2.00 moles of oxygen, and 3.00 moles of carbon dioxide at a temperature of 400 K, what is the pressure inside the cylinder?

$$n_{total} = 5.00 \text{ moles} + 2.00 \text{ moles} + 3.00 \text{ moles} = 10.00 \text{ moles total}$$

$$P_{total} = \frac{nRT}{V} = \frac{(10.00 \text{ mol})(0.0821 \text{ L}\cdot\text{atm/mol}\cdot\text{K})(400 \text{ K})}{40.0 \text{ L}} = 8.21 \text{ atm}$$

A Molecular View of the Gas Laws



Diffusion

The spread of particles through random motion.


Lighter particles diffuse more quickly.




Westend61 / Superstock

Effusion


The process of a gas escaping from a container.



Effusion, Continued



Gas Stoichiometry, Part 1




Scott Eisen/Bloomberg via Getty Images

Stoichiometry


Gas Laws

Gas Stoichiometry, Part 2



Revell, *Introductory Chemistry*,
2e, © 2021 W. H. Freeman and
Company


Gas Stoichiometry, Part 3



Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Gas Stoichiometry Practice

In the fermentation of glucose, how many moles of carbon dioxide are produced for each kilogram of glucose that reacts? If the reaction takes place in a sealed container and the gas occupies a volume of 8.10 liters at a temperature of 21 °C, find the pressure of the carbon dioxide gas inside the container.



g $\text{C}_6\text{H}_{12}\text{O}_6 \Rightarrow$ Moles $\text{C}_6\text{H}_{12}\text{O}_6 \Rightarrow$ Moles CO_2

$$1,000 \text{ g } \cancel{\text{C}_6\text{H}_{12}\text{O}_6} \times \frac{1 \text{ mol } \cancel{\text{C}_6\text{H}_{12}\text{O}_6}}{180.18 \text{ g } \cancel{\text{C}_6\text{H}_{12}\text{O}_6}} \times \frac{2 \text{ mol CO}_2}{1 \text{ mol } \cancel{\text{C}_6\text{H}_{12}\text{O}_6}} = 11.10 \text{ mol CO}_2$$


Moles $\text{CO}_2 \Rightarrow$ Pressure CO_2

$$P = \frac{nRT}{V} = \frac{(11.10 \text{ mol } \cancel{\text{CO}_2})(0.0821 \text{ L} \cdot \text{atm} / \text{mol} \cdot \cancel{\text{K}})(294 \text{ K})}{8.10 \text{ L}} = 33.1 \text{ atm CO}_2$$

$$T = 21^\circ\text{C} + 273 = 294 \text{ K}$$

Gas Stoichiometry, More Practice

Natural gas burns cleanly in air, according to this equation:



If 13.1 liters of CH_4 burn at a pressure of 1.00 atmosphere and a temperature of 290 K, what mass of carbon dioxide gas is produced?



Vitaly Maslko/Alamy

Pressure, Volume $\text{CH}_4 \Rightarrow$ Moles CH_4


$$n = \frac{PV}{RT} = \frac{(1.00 \text{ atm})(13.1 \text{ L})}{(0.0821 \text{ L}\cdot\text{atm}/\text{mol}\cdot\text{K})(290 \text{ K})} = 0.550 \text{ moles } \text{CH}_4$$

Moles $\text{CH}_4 \Rightarrow$ Moles $\text{CO}_2 \Rightarrow$ Grams CO_2

$$0.550 \text{ mol } \text{CH}_4 \times \frac{1 \text{ mol } \text{CO}_2}{1 \text{ mol } \text{CH}_4} \times \frac{44.01 \text{ g } \text{CO}_2}{1 \text{ mol } \text{CO}_2} = 24.2 \text{ g } \text{CO}_2$$

Gas Stoichiometry Summary

The mole map, including gases



Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company